

Matter in F-Theory

Yu. Malyuta and T. Obikhod

*Institute for Nuclear Research
National Academy of Sciences of Ukraine
252022 Kiev, Ukraine
e-mail: interdep@kinr.kiev.ua*

Abstract

It is shown that the matter content of F-theory compactifications on elliptic Calabi-Yau threefolds is encoded in the Gromov-Witten invariants.

1 Introduction

The matter content of six-dimensional theories resulting from compactification of F-theory on elliptic Calabi-Yau threefolds has been obtained in [1] by using Tate's algorithm.

Another algorithm for obtaining the matter content of six-dimensional theories has been proposed in [2]. This algorithm allows to obtain the matter content from the generalized Green-Schwarz anomaly cancellation condition.

The purpose of the present paper is to show that the matter content of F-theory compactifications on elliptic Calabi-Yau threefolds is encoded in the Gromov-Witten invariants.

2 Gromov-Witten invariants

Let us consider the chains of elliptic fibred Calabi-Yau threefolds recorded in Table 1, where F_n is the Hirzebruch surface and (h^{21}, h^{11}) are the Hodge numbers.

F_n	$SU(1)$	$SU(2)$	$SU(3)$	$SU(4)$	$SU(5)$	$SO(10)$	E_6	E_7
F_2	(243, 3)	(190, 4)	(161, 5)	(138, 6)	(117, 7)	(112, 8)	(105, 9)	(96, 10)
F_4	(271, 7)	(194, 8)	(153, 9)	(122, 10)	(95, 11)	(88, 12)	(79, 13)	(68, 14)
F_6	(321, 9)	(220, 10)	(167, 11)	(128, 12)	(95, 13)	(86, 14)	(75, 15)	(62, 16)
F_8	(376, 10)	(251, 11)	(186, 12)	(139, 13)	(100, 14)	(89, 15)	(76, 16)	(61, 17)
F_{10}	(433, 13)	(284, 14)	(207, 15)	(152, 16)	(107, 17)	(94, 18)	(79, 19)	(62, 20)
F_{12}	(491, 11)	(318, 12)	(229, 13)	(166, 14)	(115, 15)	(100, 16)	(83, 17)	(64, 18)

Table 1 : The chains of elliptic fibred Calabi-Yau threefolds.

The chains for each n have similar properties, therefore we will treat in some detail only the first three models of the chain $n = 2$. Application of the program INSTANTON [3] to these models gives the Gromov-Witten invariants recorded in Tables 2, 3 and 4. Note that our normalization is given by

$$\frac{1}{2}J^3,$$

where J^3 is the topological coupling [4].

(0,0,1)	-1	(0,1,1)	-1	(0,1,2)	-2	(0,1,3)	-3
(0,1,4)	-4	(0,1,5)	-5	(0,2,3)	-3	(0,2,4)	-16
(1,0,0)	240	(1,0,1)	240	(1,1,1)	240	(1,1,2)	720
(1,1,3)	1200	(1,1,4)	1680	(1,2,3)	1200	(2,0,0)	240
(2,0,2)	240	(2,2,2)	240	(3,0,0)	240	(3,0,3)	240
(4,0,0)	240	(5,0,0)	240	(6,0,0)	240	(0,1,0)	0

Table 2 : The Gromov-Witten invariants for $X_{24}(1, 1, 2, 8, 12)_{-480}^{3,243}$.

(0,0,0,1)	28	(0,0,0,2)	-1	(0,0,0,3)	0	(0,0,1,0)	-1
(0,1,0,0)	0	(0,1,1,0)	-1	(0,1,2,0)	-2	(0,1,3,0)	-3
(0,1,4,0)	-4	(0,1,5,0)	-5	(0,2,3,0)	-3	(0,2,4,0)	-16
(0,2,5,0)	-55	(0,2,6,0)	-144	(0,3,4,0)	-4	(0,3,4,0)	-4
(0,3,5,0)	-55	(1,0,0,0)	-1	(1,0,0,1)	28	(1,0,0,2)	186
(1,0,0,3)	28	(1,0,0,4)	-1				

Table 3 : The Gromov-Witten invariants for $X_{20}(1, 1, 2, 6, 10)_{-372}^{4,190}$.

(0,0,0,0,1)	15	(0,0,0,0,2)	0	(0,0,0,1,0)	-1	(0,0,0,1,1)	15
(0,0,0,1,2)	15	(0,0,0,1,3)	-1	(0,0,1,0,0)	-1	(0,1,0,0,0)	0
(0,1,1,0,0)	-1	(0,1,2,0,0)	-2	(0,1,3,0,0)	-3	(0,1,4,0,0)	-4
(0,2,3,0,0)	-3	(0,2,4,0,0)	-16	(0,2,5,0,0)	-55	(1,0,0,0,0)	-1
(1,0,0,1,0)	-1	(1,0,0,1,1)	15	(1,0,0,1,2)	15	(1,0,0,1,3)	-1
(1,0,0,2,2)	15	(1,0,0,2,3)	156	(1,0,0,2,4)	15	(1,0,1,0,0)	-1
(1,0,0,3,0)	-1	(1,0,0,3,1)	15	(1,0,0,3,2)	15	(1,0,0,3,3)	-1

Table 4 : The Gromov-Witten invariants for $X_{18}(1, 1, 2, 6, 8)_{-312}^{5,161}$.

3 Matter content

From Tables 2 and 3 we infer the relation for the Gromov-Witten invariants

$$n_{a,b,c} = \sum_k n_{a,b,c,k} . \quad (1)$$

This relation describes the phase transition between the models $X_{24}(1, 1, 2, 8, 12)_{-480}^{3,243}$ and $X_{20}(1, 1, 2, 6, 10)_{-372}^{4,190}$. Let us consider the following realization of (1)

$$240 = -1 + 28 + 186 + 28 - 1 . \quad (2)$$

The matter content of the model $X_{20}(1, 1, 2, 6, 10)_{-372}^{4,190}$ can be read off from (2). The result is

$$28 \quad \mathbf{2} .$$

From Tables 3 and 4 we infer the relation for the Gromov-Witten invariants

$$n_{a,b,c,d} = \sum_k n_{a,b,c,d,k} . \quad (3)$$

This relation describes the phase transition between the models $X_{20}(1, 1, 2, 6, 10)_{-372}^{4,190}$ and $X_{18}(1, 1, 2, 6, 8)_{-312}^{5,161}$. Let us consider the following realizations of (3)

$$\begin{aligned} 28 &= -1 + 15 + 15 -1 , \\ 186 &= 15 + 156 + 15 , \\ 28 &= -1 + 15 + 15 -1 . \end{aligned} \quad (4)$$

The matter content of the model $X_{18}(1, 1, 2, 6, 8)_{-312}^{5,161}$ can be read off from (4). The result is

$$30 \quad \mathbf{3} .$$

Applying the analogous consideration to the remaining models listed in Table 1 we obtain the matter content recorded in Table 5.

Group	Matter content
$SU(2)$	$(6n + 16)\mathbf{2}$
$SU(3)$	$(6n + 18)\mathbf{3}$
$SU(4)$	$(n + 2)\mathbf{6} + (4n + 16)\mathbf{4}$
$SU(5)$	$(3n + 16)\mathbf{5} + (2 + n)\mathbf{10}$
$SO(10)$	$(n + 4)\mathbf{16} + (n + 6)\mathbf{10}$
E_6	$(n + 6)\mathbf{27}$
E_7	$(\frac{n}{2} + 4)\mathbf{56}$

Table 5 : Matter content of models with enhanced gauge groups (for $n = 2, 4, 6, 8, 10, 12$) .

Acknowledgement

We wish to thank Professor O.S. Parasyuk for useful discussions.

References

- [1] M. Bershadsky, K. Intriligator, S. Kachru, D.R. Morrison, V. Sadov and C. Vafa, *Geometric Singularities and Enhanced Gauge Symmetries*, hep-th/9605200.
- [2] P. Candelas, E. Perevalov and G. Rajesh, *Matter from Toric Geometry*, hep-th/9707049.
- [3] S. Hosono, A. Klemm, S. Theisen and S.-T. Yau, *Mirror Symmetry, Mirror Map and Applications to Complete Intersection Calabi-Yau Spaces*, hep-th/9406055.
- [4] S. Hosono, A. Klemm, S. Theisen and S.-T. Yau, *Mirror Symmetry, Mirror Map and Applications to Calabi-Yau Hypersurfaces*, hep-th/9308122.